

AD-A033 718

STANFORD UNIV CALIF DEPT OF STATISTICS

F/0 12/1

ASYMPTOTIC PROBABILITIES IN A CRITICAL AGE-DEPENDENT BRANCHING --ETC(U)

DEC 76 H J WEINER

N00014-76-C-0475

NL

UNCLASSIFIED

TR-240

1 OF 1  
AD  
A033718




END

DATE

FILMED

2-77



ADA033718

12

DDC  
DEC 27 1976  
REGISTERED

6 ASYMPTOTIC PROBABILITIES IN A  
CRITICAL AGE-DEPENDENT BRANCHING PROCESS

by

10 Howard J. Weiner

9  
Technical Report No. 240  
December 3, 1976

14 TR-240

Prepared under Contract N00014-76-C-0475  
(NR-042-267)

15 Office of Naval Research

Herbert Solomon, Project Director

Reproduction in Whole or in Part is Permitted for  
any Purpose of the United States Government

Approved for public release; distribution unlimited

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA

11 3 Dec  
12 20 p.  
DDC  
REF  
DEC 5  
1976

by Howard J. Weiner

Let  $Z(t)$  denote the number of cells alive at time  $t$  in a critical Bellman-Harris age-dependent branching process, that is, where the mean number of offspring per parent is one. A comparison method is used to show for  $k \geq 1$ , and a high-order moment condition on  $G(t)$ , where  $G(t)$  is the cell lifetime distribution, that  $\lim_{t \rightarrow \infty} t^2 P[Z(t)=k] = a_k > 0$ , where  $\{a_k\}$  are constants.

The method is also applied to the total progeny in the critical process.

[illegible]

Asymptotic Probabilities in a  
Critical Age-Dependent Branching Process

by Howard J. Weiner

1. Introduction.

Let  $Z(t)$  denote the number of cells alive at time  $t$  in a critical Bellman-Harris [2] age-dependent branching process. That is, at time  $t = 0$ , the process starts with one new cell with lifetime distribution  $G(t)$ ,  $G(0+) = 0$ , non-lattice, and

$$(1.1) \quad G'(t) = g(t),$$

the density exists, and for some  $\delta > 0$ ,

$$(1.2) \quad \int_0^{\infty} t^{4+\delta} g(t) dt < \infty.$$

At the end of its life, the cell disappears and is immediately replaced by  $k$  new cells with probability  $p_k \geq 0$ , with

$$(1.3) \quad \sum_{k=0}^{\infty} p_k = 1$$

and criticality,

$$(1.4) \quad \sum_{k=1}^{\infty} k p_k = 1.$$

Each new cell proceeds identically as the parent cell and independently of all other cells and the state of the system. Assume that for all integers  $k$ ,

$$(1.5) \quad \sum_{n=1}^{\infty} n^k p_n < \infty.$$

It is the purpose of this note to show that for  $k \geq 1$ ,

$$(1.6) \quad \lim_{t \rightarrow \infty} t^2 P[Z(t)=k] = a_k > 0,$$

where  $\{a_k\}$  are constants.



The method is also applied to total progeny in the critical process.

## 2. Integral Equation and Result.

Define the offspring generating function, for  $0 \leq s \leq 1$ ,

$$(2.1) \quad h(s) \equiv \sum_{k=0}^{\infty} p_k s^k.$$

Define the generating function for  $Z(t)$  as, for  $0 \leq s \leq 1$ , all  $t \geq 0$ ,

$$(2.2) \quad F(s,t) \equiv E(s^{Z(t)}).$$

Then [2]  $F(s,t)$  satisfies

$$(2.3) \quad F(s,t) = 1 - G(t) + \int_0^t h(F(s,t-u)) dG(u),$$

with  $F(s,0) = s$ .

Theorem. Let  $Z(t)$  denote the number of cells at time  $t$  in a critical Bellman-Harris branching process satisfying (1.1) - (1.5). Assume also that the derivatives of  $h$  satisfy

$$(2.4) \quad 0 < h''(1) < \infty, \quad 0 < h'''(1) < \infty.$$

Then for  $k \geq 1$

$$(2.5) \quad \lim_{t \rightarrow \infty} t^2 P[Z(t)=k] = a_k > 0$$

where  $\{a_k\}$  are constants.

Proof. Consider first the case  $k = 1$ , and write for simplicity

$$(2.6) \quad P(t) \equiv P[Z(t)=1].$$

From the representation (2.3), derivatives of  $F(s,t)$  with respect to  $s$  exist, and note that

$$(2.7) \quad \left. \frac{\partial F(s,t)}{\partial s} \right|_{s=0} = P(t).$$

From (2.7), (2.3) one obtains the relationship

$$(2.8) \quad P(t) = 1 - G(t) + \int_0^t h'(1-Q(t-u))P(t-u)dG(u)$$

$$\text{and } P(0) = 1,$$

where

$$(2.9) \quad Q(t) = P[Z(t) > 0].$$

From [2],

$$(2.10) \quad \lim_{t \rightarrow \infty} tQ(t) = 2\mu(h''(1))^{-1},$$

where

$$(2.11) \quad \mu \equiv \int_0^{\infty} tdG(t).$$

Assume for simplicity in all the following that

$$(2.12) \quad \mu \equiv 1.$$



The proof will proceed by a number of claims.

Claim I. Let  $R(t)$  be continuous,  $R(0) = 0$ , and satisfy, for  $t > 0$ , that

$$(2.13) \quad R(t) \begin{matrix} > \\ (<) \end{matrix} \int_0^t h'(1-Q(t-u))R(t-u)dG(u).$$

Then for all  $t > 0$ ,

$$(2.14) \quad R(t) \begin{matrix} > \\ (<) \end{matrix} 0.$$

Proof. Assume the upper inequality of (2.13).

Note first that for all  $t > 0$ ,

$$(2.15) \quad R(t) \neq 0.$$

This follows from (2.13) by assuming there is a  $t_0$  such that  $R(t_0) = 0$ , and a contradiction is clear.

If the upper inequality in (2.13) is false, (2.15) requires that  $R(t) < 0$  for  $t > 0$ . Then on an arbitrary interval  $[0, t]$ , assume that  $R(t)$  assumes its minimum at  $t_1 \leq t$ . Since  $0 \leq h'(x) \leq 1$ , setting  $t = t_1$  in the upper inequality (2.13) yields a contradiction. The lower inequality of (2.13) implies the lower inequality of (2.14) similarly, completing the claim.

Claim II. For some  $0 < \epsilon < 1$ , all  $t \geq 0$ , define

$$(2.16) \quad K(t) = \begin{cases} 1 & , \quad t < 1 \\ \frac{1}{t^{2-\epsilon}} & , \quad t \geq 1 \end{cases}$$

$$(2.17) \quad L(t) = \begin{cases} 1 & , \quad t < 1 \\ \frac{1}{t^{2+\epsilon}} & , \quad t \geq 1. \end{cases}$$

Then for all  $t$  sufficiently large,

$$(2.18) \quad K(t) > 1 - G(t) + \int_0^t h'(1-Q(t-u))K(t-u)dG(u)$$

and

$$(2.19) \quad L(t) < 1 - G(t) + \int_0^t h'(1-Q(t-u))L(t-u)dG(u).$$

Proof. To show (2.18), note that the right side yields the inequality

$$(2.20) \quad \int_0^t h'(1-Q(t-u))K(t-u)dG(u) \leq G(t) - G(t/2) \\ + \int_0^{t/2} h'(1-Q(t-u))K(t-u)dG(u).$$

A Taylor expansion of the rhs of (2.20) yields,

$$(2.21) \quad \int_0^{t/2} h'(1-Q(t-u))K(t-u)dG(u) \\ = \int_0^{t/2} (1-Q(t-u)h''(1) + o(Q(t-u)))K(t-u)dG(u).$$

A Newton binomial expansion applied to the rhs integrand of (2.21) using (2.10), Theorem 4, p. 406 of [3], (2.16), and that  $\mu \equiv 1$  yields that the rhs of (2.21) equals

$$\begin{aligned}
(2.22) \quad & \int_0^{t/2} \left(1 - \frac{2}{t-u} + o\left(\frac{1}{t-u}\right)\right) \frac{dG(u)}{(t-u)^{2-\epsilon}} \\
&= \frac{1}{t^{2-\epsilon}} \int_0^{t/2} \frac{dG(u)}{\left(1 - \frac{u}{t}\right)^{2-\epsilon}} - \frac{2}{t^{3-\epsilon}} + o\left(\frac{1}{t^{3-\epsilon}}\right) \\
&= \frac{1}{t^{2-\epsilon}} \int_0^{t/2} \left(1 + \frac{(2-\epsilon)u}{t} + o\left(\frac{1}{t}\right)\right) dG(u) - \frac{2}{t^{3-\epsilon}} + o\left(\frac{1}{t^{3-\epsilon}}\right) \\
&= \frac{1}{t^{2-\epsilon}} + \frac{(2-\epsilon)}{t^{3-\epsilon}} - \frac{2}{t^{3-\epsilon}} + o\left(\frac{1}{t^{3-\epsilon}}\right) \\
&< \frac{1}{t^{2-\epsilon}}
\end{aligned}$$

for all  $t$  sufficiently large. In view of (2.20), (1.1), this suffices for (2.18), and (2.19) follows similarly.

Claim III. For  $0 < \epsilon < 1$ , and all  $t$  sufficiently large,

$$(2.23) \quad \frac{1}{t^{2+\epsilon}} < P(t) < \frac{1}{t^{2-\epsilon}}.$$

Proof. This follows from Claims I and II, where  $R = K - P$ , for example.

Claim IV. Let  $R(t)$  satisfy

$$(2.24) \quad R(t) = 1 - G(t) + \int_0^t h'(1-Q(t-u))R(t-u)dG_0(u),$$

with  $R(0) = 1$ , where  $Q(t)$  is the probability of non-extinction for the process with lifetime  $G(t)$ , and

$$(2.25) \quad 1 - G_0(t) = e^{-t}.$$



Then as  $t \rightarrow \infty$ ,

$$(2.26) \quad R(t) \sim \frac{c}{t^2}$$

for some constant  $c > 0$ .

Proof. Write (2.24) as

$$(2.27) \quad R(t) = 1 - G(t) + e^{-t} \int_0^t h'(1-Q(u))R(u)e^u du.$$

Differentiating, with respect to  $t$ ,

$$(2.28) \quad R'(t) = -g(t) + (1-G(t)-R(t)) + h'(1-Q(t))R(t).$$

Then, expanding  $h$  in a Taylor series, one obtains

$$(2.29) \quad R'(t) + \frac{2}{t} R(t) = f(t)$$

where

$$(2.30) \quad f(t) = 1 - G(t) - g(t) + \frac{h'''(*)}{2} Q^2(t)R(t) + (Q(t) - \frac{2}{h''(1)t})R(t),$$

with  $1 - Q(t) \leq * \leq 1$ . In view of (1.1), (1.2), (1.5), it follows from Theorem 4 of [3], p. 406, that

$$(2.31) \quad (Q(t) - \frac{2}{h''(1)t}) = O(\frac{\log t}{t^2}).$$

An analysis similar to that in Claims I - III yields that for  $0 < \epsilon < 1$ , for  $t$  sufficiently large,

$$(2.32) \quad \frac{1}{t^{2+\epsilon}} < R(t) < \frac{1}{t^{2-\epsilon}}.$$

Then (1.1), (2.31), (2.32) used in (2.30) imply that

$$(2.33) \quad t^2 f(t)$$

is integrable with respect to Lebesgue measure on the positive line.

To solve (2.29) for  $t$  large, let

$$(2.34) \quad Q(t) \equiv t^2 R(t).$$

Then (2.29) becomes

$$(2.35) \quad Q'(t) = t^2 f(t).$$

For fixed  $t_0 > 0$ , one then obtains

$$(2.36) \quad Q(t) - Q(t_0) = \int_{t_0}^t \xi^2 f(\xi) d\xi$$

or

$$(2.37) \quad R(t) \sim \frac{c}{t^2}$$

for  $t$  sufficiently large, and  $c > 0$ , completing this claim.



Claim V. Let

$$(2.38) \quad T(t) = 1 - G(t) + \int_0^t h'(1-Q(t-u))R(t-u)dG(u).$$

Then, as  $t \rightarrow \infty$ ,

$$(2.39) \quad t^2 |T(t) - R(t)| \rightarrow 0.$$

Proof.

$$(2.40) \quad |T(t) - R(t)| \leq \left| \int_0^{t/2} h'(1-Q(t-u))R(t-u)(dG(u) - dG_0(u)) \right| \\ + G(t) - G(t/2) + G_0(t) - G_0(t/2).$$

A Taylor series and Newton binomial expansion on the right side integrand of (2.40) yields

$$(2.41) \quad \int_0^{t/2} \left[ 1 - Q(t-u)h''(1) + Q^2(t-u)h'''(k(t-u)) \right] \left[ \frac{c}{(t-u)^2} + o\left(\frac{1}{(t-u)^2}\right) \right] (dG(u) - dG_0(u))$$

where

$$(2.42) \quad 1 - Q(t-u) \leq k(t-u) \leq 1,$$

and expression (2.41) equals

$$\begin{aligned}
(2.43) \quad & \int_0^{t/2} \left[ 1 - \frac{2}{t-u} + o\left(\frac{1}{t-u}\right) \right] \left[ \frac{c}{(t-u)^2} + o\left(\frac{1}{(t-u)^2}\right) \right] (dG(u) - dG_0(u)) \\
&= \int_0^{t/2} \left[ 1 - \frac{2}{t}\left(1 + \frac{u}{t}\right) + o\left(\frac{1}{t}\right) \right] \left[ \frac{c}{t^2} + o\left(\frac{1}{t^2}\right) \right] (dG(u) - dG_0(u)) \\
&= o(t^{-4}),
\end{aligned}$$

for  $t$  sufficiently large, using that

$$\mu = \int_0^\infty u dG(u) = \int_0^\infty u dG_0(u) = 1.$$

This completes Claim V.

The proof of the theorem for  $P(t)$  may now be completed.

Define the iterative scheme, for  $n \geq 0$

$$(2.44) \quad P_{(n+1)}(t) = 1 - G(t) + \int_0^t h'(1-Q(t-u)) P_{(n)}(t-u) dG(u),$$

and let

$$(2.45) \quad P_{(0)}(t) \equiv R(t).$$

Then from (2.38), it follows that

$$(2.46) \quad P_{(1)}(t) = T(t).$$

Claims IV and V imply that, for  $t$  large,

$$(2.47) \quad P_{(0)}(t) \sim \frac{c}{t^2} \quad \text{and} \quad P_{(1)}(t) \sim \frac{c}{t^2}.$$

An induction argument along the lines of Claim V establishes that,  
for large  $t$ ,

$$(2.48) \quad P_{(n)}(t) \sim \frac{c}{t^2}.$$

Note that for  $n \geq 0$ ,

$$(2.49) \quad P(t) - P_{(n+1)}(t) = \int_0^t h'(1-Q(t-u))(P(t-u) - P_{(n)}(t-u))dG(u).$$

Denote

$$(2.50) \quad \Delta_n(t) \equiv P(t) - P_{(n)}(t),$$

then from (2.49),

$$(2.51) \quad |\Delta_{n+1}(t)| < \int_0^t |\Delta_n(t-u)|dG(u) < |\Delta_0|^* G_{n+1}(t),$$

where  $*$  denotes convolution, and  $G_n(t)$  is the  $n^{\text{th}}$  convolution of  $G$  with itself.

Let  $\{X_i\}$  denote I.I.D. random variables, with distribution function  $G(t)$ .

Set

$$(2.52) \quad S_n \equiv \sum_{i=1}^n X_i.$$

Then

$$(2.53) \quad G_n(t) \equiv P[S_n \leq t] = P[S_n - n \leq t - n],$$

and if one takes  $n > t$ , by (1.2) and Chebyshev's inequality,

$$(2.54) \quad G_n(t) \leq \frac{\text{Var } S_n}{(n-t)^2} = \frac{n}{(n-t)^2}.$$

Also,

$$(2.55) \quad |\Delta_o| * G_n(t) \leq G(t) - G(t/2) + \int_0^{t/2} |P(t-u) - R(t-u)| dG_n(u)$$

and (1.2), Claims III, IV, and (2.54) yields that

$$(2.56) \quad |\Delta_o| * G_n(t) \leq \frac{Kn}{t^{2-\epsilon}(n-t)^2},$$

where  $K$  is a constant.

For fixed large  $t$ , let

$$(2.57) \quad n > [t^{1+\epsilon}].$$

This suffices for the proof of the theorem for  $P(t) \equiv P[Z(t) = 1]$ .

For  $k \geq 2$ , note that

$$(2.58) \quad k!P[Z(t)=k] \equiv k!P_k(t) = \left. \frac{\partial^k F(s,t)}{\partial s^k} \right|_{s=0}.$$

Then (2.58) applied to (2.3) for  $k = 2$  yields

$$(2.59) \quad P_2(t) = \frac{1}{2} \int_0^t h''(1-Q(t-u)) P^2(t-u) dG(u) + \int_0^t h'(1-Q(t-u)) P_2(t-u) dG(u).$$



Since the theorem is true for  $P(t)$ , then as (2.59) is of the same form as (2.8), a similar analysis as that used to establish the result for  $P(t)$  also establishes it for  $P_2(t)$ . Assume the result true for  $P_k(t)$ ,  $k \leq n$ . Applying (2.58) to (2.3) for  $k = n+1$  yields an equation of the form

$$(2.60) \quad P_{n+1}(t) = f_{n+1}(t) + \int_0^t h'(1-Q(t-u))P_{n+1}(t-u)dG(u)$$

where the induction hypothesis yields that

$$(2.61) \quad f(t) = o(t^{-4}).$$

In view of (2.61), equation (2.60) is of the same form as (2.8), and the analysis for  $P(t)$  applies to  $P_{n+1}(t)$ , yielding the theorem.

Remark. This result is known for the critical discrete-time or Galton-Watson process. See [2] Ch. 1, for example. It may be possible that a series of comparison sequences of the type given in (2.44), (2.45) relating the critical Bellman-Harris process with a corresponding critical Galton-Watson process could yield this result.

### 3. Total Progeny.

For another application of the method, let

(3.1)  $N(t)$  = total number of progeny born by  $t$  in a critical age-dependent branching process with  $\infty > h'(1) \equiv \sum_{k=1}^{\infty} kp_k > 0$ .

Denote, for  $0 \leq s \leq 1$ ,  $t \geq 0$

$$(3.2) \quad H(s, t) \equiv E(s^{N(t)}) = \sum_{k=1}^{\infty} P[N(t)=k]s^k.$$



Then (e.g. [4], p. 394)

$$(3.3) \quad H(s, t) = s \left[ 1 - G(t) + \int_0^t h(H(s, t-u)) dG(u) \right].$$

Since, for  $k \geq 1$ ,

$$(3.4) \quad \left. \frac{\partial^k H(s, t)}{\partial s^k} \right|_{s=0} = k! P[N(t)=k] \equiv k! Q_k(t)$$

one may apply (3.4) to (3.3) to obtain the limiting values of  $P[N(t)=k] \equiv Q_k(t)$ .

For  $k = 1, 2$  one obtains, since  $H(0, t) \equiv 0$ , that

$$(3.5) \quad Q_1(t) = 1 - G(t) + \int_0^t h(0) dG(u) \rightarrow p_0 \quad \text{as } t \rightarrow \infty.$$

$$(3.6) \quad Q_2(t) = \int_0^{t/2} + \int_{t/2}^t h'(0) Q_1(t-u) dG(u) \rightarrow p_0 p_1 \quad \text{as } t \rightarrow \infty.$$

Similarly, by induction and an application of Leibniz' lemma for successive differentiation, the limiting values of the  $\{Q_k(t)\}$  may be obtained. This result is implied by a representation essentially as that in [1], pp. 275-276. No assumptions on higher moments of  $G$  are required, but all moments of  $h(s)$  are needed in this approach.

## REFERENCES

- [1] ATHREYA, K. B. and KARLIN, S. (1967). Limit theorems for the split times of branching processes. Jour. Math. and Mechanics 17, 257-278.
- [2] ATHREYA, K. and NEY, P. (1972). Branching Processes. Springer-Verlag, New York.
- [3] CHOVER, J. and NEY, P. (1968). The non-linear renewal equation. J. d'Analyse Mathématique, 21, 381-413.
- [4] WEINER, H. (1966). One age-dependent branching processes. J. Appl. Prob. 3, 383-402.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 240	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ASYMPTOTIC PROBABILITIES IN A CRITICAL AGE-DEPENDENT BRANCHING PROCESS		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) HOWARD J. WEINER		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0475 ✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, California		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (NR-042-267)
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics & Probability Program Code 436 Arlington, Virginia 22217		12. REPORT DATE December 3, 1976
		13. NUMBER OF PAGES 15
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release; Distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) CRITICAL BRANCHING PROCESS ASYMPTOTIC PROBABILITY AGE DEPENDENT		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  SEE REVERSE SIDE		

DD FORM 1473  
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



Asymptotic Probabilities in a  
Critical Age-Dependent Branching Process

by Howard J. Weiner

ABSTRACT

→ Let  $Z(t)$  denote the number of cells alive at time  $t$  in a critical Bellman-Harris age-dependent branching process, that is, where the mean number of offspring per parent is one. A comparison method is used to show for  $k \geq 1$ , and a high-order moment condition on  $G(t)$ , where  $G(t)$  is the cell lifetime distribution, that  $\lim_{t \rightarrow \infty} t^2 P[Z(t)=k] = a_k > 0$ , where  $\{a_k\}$  are constants.

The method is also applied to the total progeny in the critical process. ←

\* in the limit as  $t$  approaches infinity  $t$  squared  $P[...$

